
# Monitoring Kentucky's Climate: The Kentucky Mesonet at WKU

Stuart A. Foster
State Climatologist for Kentucky
Kentucky Climate Center
Western Kentucky University

Kentucky Water Resources Board Frankfort, Kentucky August 29, 2016

# Kentucky's Weather and Climate Extremes


Warren County, May 2010



Todd County, October 2010

#### Kentucky Climate Center

- History
  - Established at Western Kentucky University in 1978
  - Recognized by the AASC as the State Climate Office for Kentucky in 2002
  - National Weather Service funding awarded to build the Kentucky Mesonet in 2006



## Climate Services Partnerships







NATIONAL WEATHER SERVICE













#### A Matter of Perspective

**Geostationary Earth Observing Platforms** 





Remote sensing often provides a bird's-eye view, while people see the world from their front porch.




# **Talking Points**

- Perspectives on Drought
- Kentucky Mesonet at WKU
- Strategic Initiatives
- Questions and Discussion




# Flash Drought!






http://droughtmonitor.unl.edu/MapsAndData/MapArchive.aspx

#### Continental Evolution of 2012 Drought



# AMJ and JAS Precipitation, 1895-2011 Western Climate Division, Kentucky





#### **Notes**

Vertical blue line represents average<sup>1</sup> AMJ precipitation.

Horizontal blue line represents average<sup>1</sup> JAS precipitation.

Vertical red line represents actual 2012 AMJ precipitation.

Dashed red line represents precipitation for JAS of 2012 required to bring the combined AMJ and JAS total to the average<sup>1</sup>. Average is defined as the arithmetic mean of the climate division values for 1895 through 2011.



#### Dimensions of the 2012 Drought



Mayfield, Graves County July 27, 2012

#### **Graves County**

- 3.22" Precipitation on March 8<sup>th</sup>
- 4.40" Precipitation for the entire spring season
- 99.4° Average high temperature from June 28<sup>th</sup> through July 8<sup>th</sup>

#### Historical Context for the Drought of 2012



Kentucky's Western Climate Division






#### Synoptic Precipitation Pattern in Late July





## Evolution of the 2012 Drought in Kentucky





#### Dimensions of Drought in Space and Time









- Onset
- Duration
- Peak Intensity
- Extent



#### Palmer Drought Severity Index



#### Types of Drought



Sequence of drought occurrence and impacts for commonly accepted drought types. All droughts originate from a deficiency of precipitation or meteorological drought but other types of drought and impacts cascade from this deficiency. (Source: National Drought Mitigation Center, University of Nebraska-Lincoln, U.S.A.) <a href="http://drought.unl.edu/DroughtBasics/TypesofDrought.aspx">http://drought.unl.edu/DroughtBasics/TypesofDrought.aspx</a>

#### Kentucky's Climatological Record

- 1825 U.S. Army
- 1861 Smithsonian Institute
- 1874 U.S. Army Signal Corps
- 1891 U.S. Department of Agriculture
- 1940 U.S. Department of Commerce
- 2007 Kentucky Climate Center



Newport Barracks, July 1825

**NWS COOP - Manual** 



**NWS COOP - Automated** 




Kentucky Mesonet







# Kentucky's diverse terrain creates distinct local vulnerabilities to weather and climate





#### Across the Commonwealth














### Vertically Integrated System









# Site Surveys and Selection

Candidate sites are surveyed and scored

Sites are selected with input from NWS and

local stakeholders

	Survey Scores	
	Temperature	40
	Precipitation	30
	Wind	12
ole Calleton	Soil	15
	Total	97



#### Station Installation and Maintenance

- Technicians install stations and instrumentation
- Technicians make spring, summer, and winter maintenance passes
- Technicians respond to "trouble tickets" when QA processes indicate problems



# Design Criteria Emphasize Quality and Reliability



**Temperature** 



Precipitation

#### Sensor Package


- Air temperature
- Precipitation
- Solar radiation
- Relative humidity
- Wind speed & direction
- Soil moisture & temperature\*

\* selected sites

Package enables ET calculations




#### Metadata Database



#### KENTUCKY MESSNET | WKU. MESONET DATA FLOW AT&T Cell Tower AT&T Custom APN Datalogger Cellular Modem **Product Generation** Mesonet Staff Ingest / Quality Assurance Server Rack Web Server **Public Safety Users Public Users** Private / Media Users Internet Consortium Users



#### **Quality Assurance Procedures**



- Automated QA runs on fiveminute data as they are collected from remote sites
- Manual QA is implemented on a daily basis to provide expert assessment of system performance




#### Meteorological Database


```
Each station collects over 105,000 observations each
         on returns over 2,730,000 data values
```

KYMN, ALBN, TBL 5min, "2010-05-01 14:55:00", "2010-05-01 09:55:00 CDT", "2010-05-01 08:55:00 CST", 62.875, 96.8, 2.55, 139.6, 4.17, 146.7, 38.07, 0.0138

#### KENTUCKY MESSNET | WKU.


The Commonwealth's Official Source for Weather and Climate Data








#### Kentucky Mesonet Coverage Map



# Kentucky Climate Center Strategic Initiatives





#### Strategic Initiatives

#### Kentucky Mesonet at WKU

Kentucky Mesonet

- Add 10-12 stations to fill coverage gaps and custom stations to meet local demand
- Enhanced instrumentation package
  - Soil probes (where not currently deployed)
  - Multi-level temperature and wind
  - Add barometric pressure sensors
- Add 3 atmospheric flux monitoring stations


Targeted completion by FY19



#### Strategic Initiatives

#### Forecasting & Outlooks

- Develop an operational unit to provide customized weather forecasts and climate outlooks
  - Hire staff meteorologist and student interns
  - Utilize existing NWS forecast model output and develop option to generate forecasts incorporating mesonet data
  - Provide client-focused forecast and outlook products. For agriculture, these would include
    - forecasts relating to stress on crops and livestock
    - outlooks for growing degree days
    - forecasts for irrigation scheduling



# Strategic Initiatives Climate Modeling

- Conduct meso-scale simulations to model impacts of land use/land cover change on atmospheric conditions
- Provide high-resolution, downscaled climate model projections for Kentucky based on model runs reported in the most recent National Climate Assessment
  - Models can be run for various time horizons

